

# Astroparticle Physics with Multiple Messengers

- Cosmic radiation from our Galaxy
- Extragalactic Cosmic Radiation
- Open Questions: Nature of the sources, chemical composition
- Role of cosmic magnetic fields
- Ultra-High Energy Cosmic Rays and secondary  $\gamma$ -rays and neutrinos: Constraints and detection prospects with different experiments.
- Testing physics beyond the Standard Model: Cross sections at PeV scales, Lorentz symmetry violation

**Günter Sigl**

**II. Institut theoretische Physik, Universität Hamburg**


**<http://www2.iap.fr/users/sigl/homepage.html>**

## The structure of the spectrum and scenarios of its origin


supernova remnants

wind supernovae



AGN, top-down ??



## Another View of the All Particle Spectrum




# Supernova Remnants and Galactic Cosmic and $\gamma$ -Rays



Aharonian et al., Nature 432 (2004) 75

Supernova remnants have been seen by HESS in  $\gamma$ -rays: The remnant RXJ1713-3946 has a spectrum  $\sim E^{-2.2}$ :  $\Rightarrow$  Charged particles have been accelerated to  $> 100$  TeV. Also seen in 1-3 keV X-rays (contour lines from ASCA)




Hadronic versus leptonic model of SN remnant HESS J1813-178:  
both are still possible




But in some supernova remnants the magnetic field needed to explain relative height of synchrotron and inverse Compton peak in the leptonic model would be too high:


$$\frac{P_{\text{synch}}}{P_{\text{IC}}} = \frac{u_B}{u_{\text{CMB+IR}}}$$



„double-humped“ spectra are also typical for AGNs



## Latest example: Lobes of Centaurus A seen by Fermi-LAT




> 200 MeV  $\gamma$ -rays

Radio observations

Abdo et al., Science Express 1184656, April 1, 2010






Low energy bump = synchrotron

high energy bump = inverse Compton on CMB in  $\sim 0.85 \mu\text{G}$  field

Abdo et al., *Science Express* 1184656, April 1, 2010

## Core of Centaurus A seen by Fermi-LAT



Can be explained by synchrotron self Compton except for HESS observation

Abdo et al., (Fermi LAT collaboration), arXiv:1006.5463

## Interactions of Hadronic primary cosmic rays

$\gamma$ -rays can be produced by  $pp \rightarrow pp\pi^0 \rightarrow pp\gamma\gamma$

$$\sigma_{pp}(s) \simeq [35.49 + 0.307 \ln^2 (s/28.94 \text{ GeV}^2)] \text{ mb}$$


This cross section is almost constant  $\rightarrow$  secondary spectra roughly the same shape as primary fluxes as long as meson cooling time is much larger than decay time.

$\gamma$ -rays can also be produced by **py interactions**:

For sub-MeV photons the cross section has a threshold and is typically  $\sim 100$  mb and weakly energy dependent at energies much above the threshold

11


$\Rightarrow$  Secondary neutrino flux also has a (very high energy) threshold above which it roughly follows the primary spectrum.





## HESS sources: X-ray binary LS 5039

Secondary  $\gamma$ -rays  
and neutrinos  
mostly produced  
by pp interactions  
in this model




F.Aharonian et al., astro-ph/0508658

Expected neutrino fluxes above TeV  $\sim 10^{-9}$ - $10^{-7}$   $\text{GeV cm}^{-2}\text{s}^{-1}$




# Hadronic Interactions and Galactic Cosmic and $\gamma$ -Rays




HESS has observed  $\gamma$ -rays from objects around the galactic centre which correlate well with the gas density in molecular clouds for a cosmic ray diffusion time of  $T \sim R^2/D \sim 3 \times 10^3 (\theta/1^\circ)^2/\eta$  years where  $D = \eta \cdot 10^{30} \text{ cm}^2/\text{s}$  is the diffusion coefficient for protons of a few TeV.


Aharonian et al., Nature 439 (2006) 695




## Identifying galactic sources from their secondary gamma-ray signatures



Shell-type supernova remnant RCW 86 seen by HESS









Given the observed spectrum  $E^{-2.3}$ , this can be interpreted as photons from  $\pi^0$  decay produced in pp interactions where the TeV protons have the same spectrum and could have been produced in a SN event.

Note that this is consistent with the source spectrum both expected from shock acceleration theory and from the cosmic ray spectrum observed in the solar neighborhood,  $E^{-2.7}$ , corrected for diffusion in the galactic magnetic field,  $j(E) \sim Q(E)/D(E)$ .

# Galactic Cosmic Ray Propagation and Signatures of Dark Matter Annihilation





## Galactic Cosmic Ray Propagation

Galactic propagation is described by solving the diffusion-convection-energy loss equation:

$$\partial_t n = \nabla \cdot (D_{xx} \nabla n - \mathbf{v}_c) + \partial_p \left( p^2 D_{pp} \partial_p \frac{n}{p^2} \right) - \partial_p \left[ \dot{p} n - \frac{p}{3} (\nabla \cdot \mathbf{v}_c n) \right] + Q(\mathbf{r}, p)$$

spatial diffusion

convection

reacceleration

energy loss


adiabatic  
compression/  
expansion

source term

This equation is solved in a cylindrical slab geometry with suitable boundary Conditions.

Out of the resulting electron/positron distribution one can compute synchrotron emission (and also inverse Compton scattering) along any line of sight.

Note: Propagation parameters are constrained by secondary to primary ratios:



Evoli, Gaggero, Grasso, Maccione, JCAP 0810, 018 (2008)

## Propagation Models

Definition of diffusion coefficients:

$$D_{xx} = \frac{v}{c_0} D_0 \left( \frac{E/Z}{\text{GV}} \right)^\delta$$

$$D_{pp} = \frac{4p^2 v_A^2}{3\delta(4 - \delta^2)(4 - \delta) D_{xx}}$$

where  $v_A$  is the Alfven speed

Models often considered:


Model	$\delta$	$D_0$ [kpc <sup>2</sup> /Myr]	$R$ [kpc]	$L$ [kpc]	$V_c$ [km/s]	$dV_c/dz$ km/s/kpc	$V_a$ [km/s]
MIN	0.85/0.85	0.0016	20	1	13.5	0	22.4
MED	0.70/0.70	0.0112	20	4	12	0	52.9
MAX	0.46/0.46	0.0765	20	15	5	0	117.6
DC	0/0.55	0.0829	30	4	0	6	0
DR	0.34/0.34	0.1823	30	4	0	0	32

# All Particle Spectrum and chemical Composition



Heavy elements start to dominate above knee

Rigidity ( $E/Z$ ) effect: combination of deconfinement and maximum energy


Hoerandel, astro-ph/0702370



3.) The knee is  
field as su





c magnetic  
SCADE:





# Do Cosmic Ray Anisotropies at 1-100 TeV reveal the Sources ?



Observed level  $\sim 10^{-3}$  is surprisingly high and difficult to explain:

wrong structure for Compton-Getting effect

too large for sources like Vela and beyond ( $> 100$  pc) because gyro-radius  $< 0.1$  pc

propagation mode, magnetic field structure ?