

SEARCH FOR NEUTRINOS FROM FLARING BLAZARS


Michael Kreter, Kerstin Fehn, Clancy James,
Thomas Eberl, Matthias Kadler, Cornelia Müller

Astroparticle School, 2014




NEUTRINOS FROM AGN

SEARCH FOR
NEUTRINOS WITH
ANTARES



Credit: S. Signoret, Univ. Arizona.



U. F. Katz, C. Spiering, arXiv: 1111.0507

NEUTRINOS FROM
AGN

MULTIMESSANGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY

MULTIMESSANGER ANALYSIS

SEARCH FOR
NEUTRINOS WITH
ANTARES

- TANAMI

NEUTRINOS FROM
AGN


MULTIMESSANGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY



SEARCH FOR NEUTRINOS WITH ANTARES


- NEUTRINOS FROM
AGN
- MULTIMESSENGER
- SOURCE SAMPLE
- ANALYSIS
- FOLLOW-UP
- MASTER THESIS
- SUMMARY



MULTI-MESSANGER ANALYSIS

SEARCH FOR
NEUTRINOS WITH
ANTARES

- ANTARES



NEUTRINOS FROM
AGN

MULTI-MESSANGER

SOURCE SAMPLE



ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY

THE ANTARES DETECTOR

- located in 2500 m depth
- 12 vertical detection lines (a ≈ 450 m)
- 885 optical modules (OMs)
- 25 storeys per line (a 3 OMs)



SEARCH FOR
NEUTRINOS WITH
ANTARES

NEUTRINOS FROM
AGN

MULTIMESSENGER


SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY

DETECTION PRINCIPLE



SEARCH FOR
NEUTRINOS WITH
ANTARES

NEUTRINOS FROM
AGN


MULTIMESSENGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY



Sky in equatorial coordinates:

- TANAMI sources
- ★ selected sources for this analysis

NEUTRINOS FROM
AGN


MULTIMESSENGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY



Selection criteria:

$$\frac{F_{\max}}{\text{meanflux}} \cdot \frac{1}{n} \quad (1)$$

NEUTRINOS FROM
AGN

MULTIMESSENGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY

SOURCE SAMPLE

SEARCH FOR
NEUTRINOS WITH
ANTARES

NEUTRINOS FROM
AGN


MULTIMESSENGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY




Selection criteria:

$$\frac{F_{\max}}{\text{meanflux}} \cdot \frac{1}{n} \quad (2)$$

AIM OF THE ANALYSIS

Enhance sensitivity in finding neutrinos from AGN in a stacked search combining

- angle
- energy
- time information



SEARCH FOR
NEUTRINOS WITH
ANTARES

NEUTRINOS FROM
AGN

MULTIMESSENGER

SOURCE SAMPLE

ANALYSIS



FOLLOW-UP
MASTER THESIS

SUMMARY

Maximize the likelihood of the data

$$L(n_s) = \prod_{i=1}^N \left[\frac{n_s}{N} S_i + \left(1 - \frac{n_s}{N}\right) B_i \right] \quad (3)$$

- n_s unknown contribution of the signal events
- N number of events
- S_i signal probability density
- B_i background probability density



NEUTRINOS FROM
AGN



MULTIMESSENGER

SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY



NEUTRINOS FROM
AGN

MULTIMESSENGER

SOURCE SAMPLE


ANALYSIS

FOLLOW-UP
MASTER THESIS

SUMMARY

Problem:

- Highly variable background from environmental conditions.




⇒ optimize MC for every single run (rbr MC)

Aim:


- Increase sensitivity by taking duration of run into account?

Idea:


- weight every neutrino event by the duration of its run



Result:



Result:



FOLLOW-UP MASTER THESIS

SEARCH FOR
NEUTRINOS WITH
ANTARES

NEUTRINOS FROM
AGN

MULTIMESSANGER


SOURCE SAMPLE

ANALYSIS

FOLLOW-UP
MASTER THESIS


SUMMARY


Result:



- Time correlation between γ and ν increases the sensitivity dramatically
- For rbr MC analysis no difference in sensitivity could be found for highly ranked sources.
- Sensitivity shift is due to bad detector contributions and high noise rates.

Backup

		signal event	background event
expectation μ		0.5, 1.0, 1.5, ..., 10.0	
S-term	angle	<div> <div>from E^{-1} MC</div> <div>(angle1, angle2, ...)</div> <div>height</div> </div>	$\sim \sin \alpha$
	energy	<div> <div>from E^{-1} MC</div> <div>(energy1, energy2, ...)</div> <div>height</div> </div>	<div> <div>from $E^{-3.6}$ MC</div> <div>height</div> <div>from E^{-1} MC</div> </div>
	time	from cumulated <u>lightcurve</u>	random
B-term		<div> <div>from E^{-1} MC</div> <div>$\frac{1}{5^{0.2} \pi}$ energy</div> <div>height</div> </div>	<div> <div>from $E^{-3.6}$ MC</div> <div>$\frac{1}{5^{0.2} \pi}$ energy</div> <div>height</div> </div>



Sources are weighted according to:

- select sources with $\frac{F_{\max}}{\text{meanflux}} > 5$
- Count the number n of bins $\text{flux} > \text{meanflux} + 3\sigma$
- ranking:

$$\frac{F_{\max}}{\text{meanflux}} \cdot \frac{1}{n}$$